



Complete Genome Sequence of *Bacillus altitudinis* P-10, a Potential Bioprotectant against *Xanthomonas oryzae* pv. *oryzae*, Isolated from Rice Rhizosphere in Java, Indonesia

Anto Budiharjo,^a Haeyoung Jeong,^b Dyah Wulandari,^c Soohyun Lee,^b Choong-Min Ryu^b

Biology Department, Faculty of Sciences and Mathematics, Diponegoro University, Semarang, Indonesia^a; Infectious Disease Research Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Daejeon, Republic of Korea^b; School of Biotechnology, Institute of Agricultural Technology, Suranaree University of Technology, Ratchasima, Thailand^c

ABSTRACT *Bacillus altitudinis* P-10 was isolated from the rhizosphere of rice grown in an organic rice field and provides strong antagonism against the bacterial blight caused by *Xanthomonas oryzae* pv. *oryzae* in rice. Herein, we provide the complete genome sequence and a possible explanation of the antibiotic function of the P-10 strain.

The species *Bacillus altitudinis* and three other novel species (*B. aerius*, *B. aerophilus*, and *B. stratosphericus*) were isolated and characterized from air samples in 2006 (1). This species was later recognized as an enzyme producer with potential industrial uses (2–6) and as a plant growth-promoting rhizobacterium (7). In addition, endophytic *B. altitudinis* isolated from the roots of healthy wheat showed the ability to bioleach iron, an important feature for lightening iron toxicity in wheat (8). However, because its 16S rRNA gene sequence is highly similar to those of *B. pumilus* group species, such as *B. pumilus*, *B. safensis*, *B. stratosphericus*, *B. altitudinis*, and *B. aerophilus* (9), a multilocus sequence analysis or a genome sequence-based analysis is required for a study of the diversity and phylogeny of this group.

B. altitudinis strain P-10 was isolated from the rhizosphere of rice in an organic rice farm in the Banyubiru District, Semarang Regency, central Java Province, Indonesia (10). Instead of a chemical fertilizer, a microbe-based biofertilizer has been applied to this rice field for a long period of time. The farmers use a biofertilizer solely to cultivate crops on this farm, because the soil, generated from an ash component created through volcanic activity, does not contain enough nutrients for bacterial growth. *B. altitudinis* strain P-10 displays direct antagonism against *Xanthomonas oryzae* pv. *oryzae*, a casual pathogen of rice bacterial blight, which has caused significant yield loss in Asian counties (10).

Genome sequencing was carried out on a PacBio RSII platform from a single-molecule real-time (SMRT) cell using P6-C4 chemistry at the Theragen Etx Bio Institute (Suwon, Gyeonggi-do, Republic of Korea). *De novo* assembly using the RS_HGAP_Assembly.2 protocol under SMRT Analysis 2.3.0 produced one contig from 1.27 Gb of raw sequencing reads, followed by circularization using Circlator (11). Residual errors were corrected by running two successive rounds of the RS_Resequencing.1 protocol, followed by additional correction through the mapping of corrected long reads in the CLC Genomics Workbench (Qiagen Bioinformatics, Aarhus, Denmark). The complete genome sequence was annotated using NCBI's Prokaryotic Genome Annotation Pipe-

Received 4 November 2017 Accepted 6 November 2017 Published 30 November 2017

Citation Budiharjo A, Jeong H, Wulandari D, Lee S, Ryu C-M. 2017. Complete genome sequence of *Bacillus altitudinis* P-10, a potential bioprotectant against *Xanthomonas oryzae* pv. *oryzae*, isolated from rice rhizosphere in Java, Indonesia. *Genome Announc* 5:e01388-17. <https://doi.org/10.1128/genomeA.01388-17>.

Copyright © 2017 Budiharjo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Choong-Min Ryu, cmryu@kribb.re.kr.

A.B. and H.J. contributed equally to this work.

line and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST) server (12).

The P-10 genome has a 3,765,752-bp chromosome (41.5% G+C content) containing 3,828 protein-coding genes, 81 tRNA genes, and 8 rRNA operons, with no plasmids. Although initially classified as *B. pumilus* based on the 16S rRNA gene sequence analysis, the Genome-to-Genome Distance Calculator (<http://ggdc.dsmz.de/ggdc.php#>) indicates that strain P-10 belongs to the *B. altitudinis* species, where the DNA-DNA hybridization (DDH) estimate with *B. altitudinis* 41KF2b^T (GenBank accession number ASJC00000000) is 86.4%.

antiSMASH 3.0 (13) predicted six biosynthetic gene clusters for lichenysin- or bacilysin-like nonribosomal peptides, terpenes, and bacteriocin. Because no notable pathogenicity-related genes have been found from Island Viewer analysis (14), *B. altitudinis* strain P-10 can be considered a promising candidate for bioprotectant development. In conclusion, the complete genome sequence of strain P-10 presents a beneficial trait and a possible explanation of its antibiotic function.

Accession number(s). The complete genome sequence of *B. altitudinis* P-10 has been deposited in DDBJ/ENA/GenBank under the accession number [CP024204](https://doi.org/10.1093/nar/gkx343).

ACKNOWLEDGMENTS

This work was supported by the KRIBB Research Initiative Program, the Advanced Biomass R&D Center (ABC) of the Global Frontier Project, funded by the Ministry of Science and ICT (grant ABC-2015M3A6A2065697), Republic of Korea, a PNBP grant from Diponegoro University, Indonesia, and the Agenda Project (PJ012814) of the Rural Development Administration (RDA).

REFERENCES

- Shivaji S, Chaturvedi P, Suresh K, Reddy GS, Dutt CB, Wainwright M, Narlikar JV, Bhargava PM. 2006. *Bacillus aerius* sp. nov., *Bacillus aerophilus* sp. nov., *Bacillus stratosphericus* sp. nov. and *Bacillus altitudinis* sp. nov., isolated from cryogenic tubes used for collecting air samples from high altitudes. *Int J Syst Evol Microbiol* 56:1465–1473. <https://doi.org/10.1099/ijs.0.64029-0>.
- Vijay Kumar E, Srijana M, Kiran Kumar K, Harikrishna N, Reddy G. 2011. A novel serine alkaline protease from *Bacillus altitudinis* GVC11 and its application as a dehairing agent. *Bioprocess Biosyst Eng* 34:403–409. <https://doi.org/10.1007/s00449-010-0483-x>.
- Esakkiraj P, Usha R, Palavesam A, Immanuel G. 2012. Solid-state production of esterase using fish processing wastes by *Bacillus altitudinis* AP-MSU. *Food Bioprod Process* 90:370–376. <https://doi.org/10.1016/j.fbp.2011.12.008>.
- Mao S, Lu Z, Zhang C, Lu F, Bie X. 2013. Purification, characterization, and heterologous expression of a thermostable beta-1, 3–1,4-glucanase from *Bacillus altitudinis* YC-9. *Appl Biochem Biotechnol* 169:960–975.
- Elbanna K, Elnaggar S, Bakeer A. 2014. Characterization of *Bacillus altitudinis* as a new causative agent of bacterial soft rot. *J Phytopathol* 162:712–722. <https://doi.org/10.1111/jph.12250>.
- Shah Mahmud R, Ulyanova V, Malanin S, Dudkina E, Vershina V, Ilinskaya O. 2015. Draft whole-genome sequence of *Bacillus altitudinis* strain B-388, a producer of extracellular RNase. *Genome Announc* 3:e01502-14. <https://doi.org/10.1128/genomeA.01502-14>.
- Sunar K, Dey P, Chakraborty U, Chakraborty B. 2015. Biocontrol efficacy and plant growth promoting activity of *Bacillus altitudinis* isolated from Darjeeling hills, India. *J Basic Microbiol* 55:91–104. <https://doi.org/10.1002/jobm.201300227>.
- Sun Z, Liu K, Zhang J, Zhang Y, Xu K, Yu D, Wang J, Hu L, Chen L, Li C-W. 2017. IAA producing *Bacillus altitudinis* alleviates iron stress in *Triticum aestivum* L. seedling by both bioleaching of iron and up-regulation of genes encoding ferritins. *Plant Soil* 419:1–11. <https://doi.org/10.1007/s11104-017-3218-9>.
- Liu Y, Lai Q, Dong C, Sun F, Wang L, Li G, Shao Z. 2013. Phylogenetic diversity of the *Bacillus pumilus* group and the marine ecotype revealed by multilocus sequence analysis. *PLoS One* 8:e80097. <https://doi.org/10.1371/journal.pone.0080097>.
- Sumarno M, Budiharjo A, Pujiyanto S. 2014. Potency and molecular identification of endospore-forming rhizobacteria from rice plant as a biocontrol of phytopathogen *Xanthomonas oryzae*, p 145–148. In Widiyandari H, Putro SP (ed), *Proceeding of the 4th International Seminar of New Paradigm and Innovation on Natural Sciences and its Application*, Diponegoro University, Semarang, Indonesia.
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.
- Bertelli C, Laird MR, Williams KP, Simon Fraser University Research, Lau BY, Hoard G, Winsor GL, Brinkman FS. 2017. IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Res* 45:W30–W35. <https://doi.org/10.1093/nar/gkx343>.